


HOW TO MIX

BRAVECTO[®]
injectable



The package includes:

- Solvent vial (1)
- Fluralaner powder vial (2)
- Vent needle (25 G) (3)
- Package leaflet


You will need to provide:

- Sterile transfer needle (18 G) (4)
- Sterile 20 ml syringe for transfer (5)

To ensure effective treatment, it is important to correctly mix the suspension before injection and use the recommended syringe and needle sizes as listed under the directions in the product information.



Begin by shaking the fluralaner powder vial to break up any aggregates prior to reconstitution.



Invert the solvent vial at least 3 times until visibly uniform. The content of the solvent vial is a clear to opaque viscous solution.

Instructions continued on next page...

HOW TO MIX BRAVECTO® injectable


3



Remove the purple cap of the sterile solvent vial. Using the 18 G needle and the 20 ml syringe, inject up to 14 ml of air into the solvent vial.


Then, withdraw 15 ml of solvent from the vial. It is important to note that there is more solvent supplied in the vial than required for reconstitution.

4




Remove the red cap of the fluralaner powder vial. **Insert a 25 gauge vent needle** into the top of this vial.

5




While **rotating the vial horizontally in your hand**, slowly transfer the solvent into the fluralaner powder vial to ensure complete wetting of the powder.

6



Once the solvent has been added, remove the vent and transfer needles from the fluralaner powder vial and discard needles.

7



Shake the vial vigorously for at least 30 seconds until a thoroughly mixed suspension of the fluralaner powder and solvent is produced. **Before every use, shake the reconstituted vial vigorously for at least 30 seconds to achieve a uniform suspension.**

BRAVECTO injectable may be stored below 30°C for 3 months after reconstitution. Discard the reconstituted vial after 20 punctures to the stopper or if the product has been suspended for more than 3 months.

For more details, view the mixing video: uk.bravovets.com



BRAVECTO® injectable (BRAVECTO® 150 mg/ml powder and solvent for suspension for injection for dogs) has a 12-months duration of efficacy against flea and tick species indicated on the SPC and contains fluralaner. **POM-V.**

Further advice should be sought from the medicine prescriber. For more information, refer to the SPC, datasheet or package leaflet.

Prescription decisions are for the person issuing the prescription alone.

Use medicines responsibly.

MSD Animal Health UK Limited. Registered office: Walton Manor, Walton, Milton Keynes MK7 7AJ, UK. Registered in England & Wales no. 946942.

© 2024 MSD Animal Health UK Limited.

UK-BRV-240300006